

Literatur

- Achttien, R., van Lieshout, J., Wensing, M., van der Sanden, M. N., & Staal, J. B. (2019). Symptoms of depression are associated with physical inactivity but not modified by gender or the presence of a cardiovascular disease; a cross-sectional study. *BMC Cardiovascular Disorders*, 19(1), 95. <https://doi.org/10.1186/s12872-019-1065-8>
- Aylett, E., Small, N., & Bower, P. (2018). Exercise in the treatment of clinical anxiety in general practice – a systematic review and meta-analysis. *BMC Health Services Research*, 18(1), 559. <https://doi.org/10.1186/s12913-018-3313-5>
- Banno M, Harada Y, Taniguchi M, et al. (2018). Exercise can improve sleep quality: a systematic review and meta-analysis. *PeerJ*, 6, e5172.
- Bellón JA. Exercise for the treatment of depression. *BMJ*. 2024 Feb 14;384:q320. doi: 10.1136/bmj.q320. PMID: 38355168.
- Birnbaum HG, Kessler RC, Kelley D, Ben-Hamadi R, Joish VN, Greenberg PE. Employer burden of mild, moderate, and severe major depressive disorder: mental health services utilization and costs, and work performance. *Depress Anxiety*. 2010;27(1):78-89. doi: 10.1002/da.20580. PMID: 19569060.
- Brinsley, J., Schuch, F., Lederman, O., Girard, D., Smout, M., Immink, M. A., Stubbs, B., Firth, J., Davison, K., & Rosenbaum, S. (2021). Effects of yoga on depressive symptoms in people with mental disorders: A systematic review and meta-analysis. *British Journal of Sports Medicine*, 55(17), 992–1000. <https://doi.org/10.1136/bjsports-2019-101242>
- Bull, F. C., Al-Ansari, S. S., Biddle, S., Borodulin, K., Buman, M. P., Cardon, G., Carty, C., Chaput, J.-P., Chastin, S., Chou, R., Dempsey, P. C., DiPietro, L., Ekelund, U., Firth, J., Friedenreich, C. M., Garcia, L., Gichu, M., Jago, R., Katzmarzyk, P. T., ... Willumsen, J. F. (2020). World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *British Journal of Sports Medicine*, 54(24), 1451–1462. <https://doi.org/10.1136/bjsports-2020-102955>

Bundesärztekammer, Kassenärztliche Bundesvereinigung, Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften, NVL Nationale Versorgungsleitlinien, Unipolare Depressionen, Version: 3.0, URL: https://www.dgppn.de/_Resources/Persistent/660e0e633be6f5db59e8f31ea3b144c29447b9b9/nvl-005k_S3_Unipolare_Depression_2022-10.pdf, (Stand 24.01.24)

Bundespsychotherapeutenkammer, Wartezeiten für psychisch kranke Menschen weiterhin zu lang: BPtK zumG-BA-Gutachten zur Bedarfsplanung (2018), URL: <https://www.bptk.de/wartezeitenfuer-psychisch-kranke-menschen-weiterhinzu-lang/>, (Stand 30.03.2020)

Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical Activity, Exercise, and Physical Fitness: Definitions and Distinctions for Health-Related Research. *Public Health Reports*, 100(2), 126–131.

Cooney, G. M., Dwan, K., Greig, C. A., Lawlor, D. A., Rimer, J., Waugh, F. R., McMurdo, M., & Mead, G. E. (2013). Exercise for depression. *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.CD004366.pub6>

Cramer, H., Lauche, R., Langhorst, J., & Dobos, G. (2013). Yoga for depression: A systematic review and meta-analysis. *Depression and Anxiety*, 30(11), 1068–1083.

DAK Gesundheit, Psychreport 2023 Entwicklungen der psychischen Erkrankungen im Job 2012-2022, Erneuter Höchststand bei psychisch bedingten Fehltagen im Job (2023), URL:<https://caas.content.dak.de/caas/v1/media/32628/data/3983614e98a936fe7d7dd70f3dac2e73/dak-psychreport-ergebnis-praesentation.pdf>, Stand (25.01.2024)

Dauwan, M., Begemann, M. J. H., Heringa, S. M., & Sommer, I. E. (2016). Exercise Improves Clinical Symptoms, Quality of Life, Global Functioning, and Depression in Schizophrenia: A Systematic Review and Meta-analysis. *Schizophrenia Bulletin*, 42(3), 588–599. <https://doi.org/10.1093/schbul/sbv164>

de Maat SM, Dekker J, Schoevers RA, de Jonghe F. Relative efficacy of psychotherapy and combined therapy in the treatment of depression: a meta-analysis. *Eur Psychiatry*. 2007 Jan;22(1):1-8. doi: 10.1016/j.eurpsy.2006.10.008. Epub 2006 Dec 27. PMID: 17194571.

Eberhard Karls Universität Tübingen, ImPuls. Starke Psyche durch Motivation und Bewegung (2023), URL: <https://www.impuls.uni-tuebingen.de/>, Stand(25.01.2024)

Ekkekakis, P. (2015). Honey, I shrunk the pooled SMD! Guide to critical appraisal of systematic reviews and meta-analyses using the Cochrane review on exercise for depression as example. *Mental Health and Physical Activity*, 8, 21–36.
<https://doi.org/10.1016/j.mhpa.2014.12.001>

GBD 2015 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016 Oct 8;388(10053):1545-1602. doi: 10.1016/S0140-6736(16)31678-6. Erratum in: *Lancet*. 2017 Jan 7;389(10064):e1. PMID: 27733282; PMCID: PMC5055577.

Guthold, R., Stevens, G. A., Riley, L. M., & Bull, F. C. (2018). Worldwide trends in insufficient physical activity from 2001 to 2016: A pooled analysis of 358 population-based surveys with 1·9 million participants. *The Lancet Global Health*, 6(10), e1077–e1086.
[https://doi.org/10.1016/S2214-109X\(18\)30357-7](https://doi.org/10.1016/S2214-109X(18)30357-7)

Guthold, R., Stevens, G. A., Riley, L. M., & Bull, F. C. (2020). Global trends in insufficient physical activity among adolescents: A pooled analysis of 298 population-based

surveys with 1·6 million participants. *The Lancet Child & Adolescent Health*, 4(1),

23–35. [https://doi.org/10.1016/S2352-4642\(19\)30323-2](https://doi.org/10.1016/S2352-4642(19)30323-2)

Heinzel, S., Lawrence, J. B., Kallies, G., Rapp, M. A., & Heissel, A. (2015). Using exercise to fight depression in older adults. *GeroPsych*.

Heissel, A., Zech, P., Rapp, M. A., Schuch, F. B., Lawrence, J. B., Kangas, M., & Heinzel, S. (2019). Effects of exercise on depression and anxiety in persons living with HIV: A meta-analysis. *Journal of Psychosomatic Research*, 126, 109823.

Jacobi F, Höfler M, Strehle J, Mack S, Gerschler A, Scholl L, Busch MA, Maske U, Hapke U, Gaebel W, Maier W, Wagner M, Zielasek J, Wittchen HU. Erratum zu: Psychische Störungen in der Allgemeinbevölkerung. Studie zur Gesundheit Erwachsener in Deutschland und ihr Zusatzmodul "Psychische Gesundheit" (DEGS1-MH) [Erratum to: Mental disorders in the general population. Study on the health of adults in Germany and the additional module mental health (DEGS1-MH)]. *Nervenarzt*. 2016 Jan;87(1):88-90. German. doi: 10.1007/s00115-015-4458-7. Erratum for: *Nervenarzt*. 2014 Jan;85(1):77-87. PMID: 26601984.

Kammerer, K., Falk, K., Heintze, C., Döpfmer, S., & Heusinger, J. (2019). Hindernisse und Voraussetzungen für die Vermittlung älterer, depressiv erkrankter Menschen in Psychotherapie aus Sicht von HausärztlInnen. *Das Gesundheitswesen*, 81(01), 58–62. <https://doi.org/10.1055/s-0042-116589>

Kendra MS, Mohr JJ, Pollard JW. The stigma of having psychological problems: relations with engagement, working alliance, and depression in psychotherapy. *Psychotherapy* (Chic). 2014 Dec;51(4):563-573. doi: 10.1037/a0036586. Epub 2014 May 26. PMID: 24866971.

LeardMann, C. A., Kelton, M. L., Smith, B., Littman, A. J., Boyko, E. J., Wells, T. S., Smith, T. C., & For the Millennium Cohort Study Team. (2011). Prospectively Assessed

Posttraumatic Stress Disorder and Associated Physical Activity. *Public Health Reports*, 126(3), 371–383. <https://doi.org/10.1177/003335491112600311>

Lee, J., Gierc, M., Vila-Rodriguez, F., Puterman, E., & Faulkner, G. (2021). Efficacy of exercise combined with standard treatment for depression compared to standard treatment alone: A systematic review and meta-analysis of randomized controlled trials. *Journal of Affective Disorders*, 295, 1494–1511. <https://doi.org/10.1016/j.jad.2021.09.043>

Lingineni, R. K., Biswas, S., Ahmad, N., Jackson, B. E., Bae, S., & Singh, K. P. (2012). Factors associated with attention deficit/hyperactivity disorder among US children: Results from a national survey. *BMC Pediatrics*, 12(1), 50.

<https://doi.org/10.1186/1471-2431-12-50>

Maljanen T, Knekt P, Lindfors O, Virtala E, Tillman P, Häkkinen T; Helsinki Psychotherapy Study Group. The cost-effectiveness of short-term and long-term psychotherapy in the treatment of depressive and anxiety disorders during a 5-year follow-up. *J Affect Disord*. 2016 Jan 15;190:254-263. doi: 10.1016/j.jad.2015.09.065. Epub 2015 Oct 24. PMID: 26540079.

Mauz E, Walther L, Junker S, Kersjes C, Damerow S, Eicher S, Hölling H, Müters S, Peitz D, Schnitzer S, Thom J. Time trends in mental health indicators in Germany's adult population before and during the COVID-19 pandemic. *Front Public Health*. 2023 Feb 23;11:1065938. doi: 10.3389/fpubh.2023.1065938. PMID: 36908429; PMCID: PMC9995751.

Miko, H.-C., Zillmann, N., Ring-Dimitriou, S., Dorner, T. E., Titze, S., & Bauer, R. (2020). Auswirkungen von Bewegung auf die Gesundheit. *Das Gesundheitswesen*, 82(S 03), S184–S195. <https://doi.org/10.1055/a-1217-0549>

Morres, I. D., Hatzigeorgiadis, A., Stathi, A., Comoutos, N., Arpin-Cribbie, C., Krommidas, C., & Theodorakis, Y. (2019). Aerobic exercise for adult patients with major depressive

disorder in mental health services: A systematic review and meta-analysis.

Depression and Anxiety, 36(1), 39–53.

National Institute for Clinical Excellence (NICE). (2009). *Depression: The treatment and*

management of depression in adults (update). Draft for consultation. Retrieved

February 22, 2021, <https://www.nice.org.uk/guidance/cg90/documents/depression-in-adults-update-draft-nice-guideline-for-consultation2>

Nübling, Rüdiger & T, Bär & Jeschke, Karin & M, Ochs & Sarubin, Nina & Juergen,

Schmidt. (2014). Versorgung psychisch kranker Erwachsener in Deutschland -

Bedarf und Inanspruchnahme sowie Effektivität und Effizienz von Psychotherapie.

Psychotherapeutenjournal. 13. 389-397.

Panza, G. A., Taylor, B. A., MacDonald, H. V., Johnson, B. T., Zaleski, A. L., Livingston,

J., Thompson, P. D., & Pescatello, L. S. (2018). Can Exercise Improve Cognitive

Symptoms of Alzheimer's Disease? *Journal of the American Geriatrics Society*,

66(3), 487–495. <https://doi.org/10.1111/jgs.15241>

Recchia, F., Leung, C. K., Chin, E. C., Fong, D. Y., Montero, D., Cheng, C. P., Yau, S. Y., &

Siu, P. M. (2022). Comparative effectiveness of exercise, antidepressants and their

combination in treating non-severe depression: A systematic review and network

meta-analysis of randomised controlled trials. *British Journal of Sports Medicine*,

56(23), 1375–1380. <https://doi.org/10.1136/bjsports-2022-105964>

Rethorst, C. D., Wipfli, B. M., & Landers, D. M. (2009). The Antidepressive Effects of Exercise:

A Meta-Analysis of Randomized Trials. *Sports Medicine*, 39(6), 491–511.

<https://doi.org/10.2165/00007256-200939060-00004>

Rosenbaum, S., Vancampfort, D., Steel, Z., Newby, J., Ward, P. B., & Stubbs, B. (2015).

Physical activity in the treatment of post-traumatic stress disorder: a systematic

review and meta-analysis. *Psychiatry research*, 230(2), 130-136.

Schneider, F., Härter, M., & Schorr, S. (Eds.). (2017). *S3-Leitlinie/Nationale VersorgungsLeitlinie Unipolare Depression*. Springer Berlin Heidelberg.
<https://doi.org/10.1007/978-3-662-52906-5>

Schuch, F. B., Vancampfort, D., Firth, J., Rosenbaum, S., Ward, P. B., Silva, E. S., Hallgren, M., Ponce De Leon, A., Dunn, A. L., Deslandes, A. C., Fleck, M. P., Carvalho, A. F., & Stubbs, B. (2018). Physical Activity and Incident Depression: A Meta-Analysis of Prospective Cohort Studies. *American Journal of Psychiatry*, 175(7), 631–648.
<https://doi.org/10.1176/appi.ajp.2018.17111194>

Schuch, F. B., Vancampfort, D., Richards, J., Rosenbaum, S., Ward, P. B., & Stubbs, B. (2016). Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. *Journal of Psychiatric Research*, 77, 42–51. URL:
<https://doi.org/10.1016/j.jpsychires.2016.02.023>

Schuch, F. B., Silva, E. S., Hallgren, M., & Fleck, M. P. (2018). Physical Activity and Incident Depression: A Meta-Analysis of Prospective Cohort Studies. *Am J Psychiatry*, 175(7), 631–648. <https://doi.org/10.1176/appi.ajp.2018.17111194>

Schuch, F. B., Stubbs, B., Meyer, J., Heissel, A., Zech, P., Vancampfort, D., Rosenbaum, S., Deenik, J., Firth, J., Ward, P. B., Carvalho, A. F., & Hiles, S. A. (2019). Physical activity protects from incident anxiety: A meta-analysis of prospective cohort studies. *Depression and Anxiety*, 36(9), 846–858. <https://doi.org/10.1002/da.22915>

Schlack R, Hapke U, Maske U, Busch M, Cohrs S. Häufigkeit und Verteilung von Schlafproblemen und Insomnie in der deutschen Erwachsenenbevölkerung: Ergebnisse der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1) [Frequency and distribution of sleep problems and insomnia in the adult population in Germany: results of the German Health Interview and Examination Survey for Adults (DEGS1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2013 May;56(5-6):740-8. German. doi: 10.1007/s00103-013-1689-2. PMID: 23703493.

Shinohara K, Honyashiki M, Imai H, Hunot V, Caldwell DM, Davies P, Moore TH, Furukawa TA, Churchill R. Behavioural therapies versus other psychological therapies for depression. *Cochrane Database Syst Rev*. 2013 Oct 16;2013(10):CD008696. doi: 10.1002/14651858.CD008696.pub2. PMID: 24129886; PMCID: PMC7433301.

Singh, B., Olds, T., Curtis, R., Dumuid, D., Virgara, R., Watson, A., Szeto, K., O'Connor, E., Ferguson, T., Eglitis, E., Miatke, A., Simpson, C. E., & Maher, C. (2023). Effectiveness of physical activity interventions for improving depression, anxiety and distress: An overview of systematic reviews. *British Journal of Sports Medicine*, 57(18), 1203–1209. <https://doi.org/10.1136/bjsports-2022-106195>

Sormunen, E., Saarinen, M. M., Salokangas, R. K. R., Telama, R., Hutri-Kähönen, N., Tammelin, T., Viikari, J., Raitakari, O., & Hietala, J. (2017). Effects of childhood and adolescence physical activity patterns on psychosis risk—A general population cohort study. *Npj Schizophrenia*, 3(1), 5. <https://doi.org/10.1038/s41537-016-0007-z>

Spörndly-Nees, S., Åsenlöf, P., & Lindberg, E. (2017). High or increasing levels of physical activity protect women from future insomnia. *Sleep Medicine*, 32, 22–27.

<https://doi.org/10.1016/j.sleep.2016.03.017>

Statistisches Bundesamt, Krankheitskosten pro Kopf gleichen sich zwischen Männern und Frauen weiter an (2022), URL:

https://www.destatis.de/DE/Presse/Pressemitteilungen/2022/07/PD22_316_236.html zugegriffen am 25.01.2024

Stubbs, B., Vancampfort, D., Rosenbaum, S., Firth, J., Cosco, T., Veronese, N., Salum, G. A., & Schuch, F. B. (2017). An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: A meta-analysis. *Psychiatry Research*, 249, 102–108. <https://doi.org/10.1016/j.psychres.2016.12.020>

Wittchen HU, Jacobi F, Rehm J, Gustavsson A, Svensson M, Jönsson B, Olesen J, Allgulander C, Alonso J, Faravelli C, Fratiglioni L, Jennnum P, Lieb R, Maercker A, van Os J, Preisig

M, Salvador-Carulla L, Simon R, Steinhausen HC. The size and burden of mental disorders and other disorders of the brain in Europe 2010. *Eur Neuropsychopharmacol.* 2011 Sep;21(9):655-79. doi: 10.1016/j.euroneuro.2011.07.018. PMID: 21896369.

Wolf, S., Seiffer, B., Zeibig, J.-M., Frei, A. K., Studnitz, T., Welkerling, J., Meinzinger, E., Bauer, L. L., Baur, J., Rosenstiel, S., Fiedler, D. V., Helmhold, F., Ray, A., Herzog, E., Takano, K., Nakagawa, T., Günak, M. M., Kropp, S., Peters, S., ... Ehring, T. (2023). A transdiagnostic group exercise intervention for mental health outpatients in Germany (ImPuls): results of a pragmatic, multi-site, block-randomised, phase 3 controlled trial. *MedRxiv*, 2023.12.16.23300028.
<https://doi.org/10.1101/2023.12.16.23300028>

Wolf, Sebastian & Zeibig, Johanna-Marie & Hautzinger, Martin & Sudeck, Gorden. (2020). Psychische Gesundheit durch Bewegung - ImPuls. Ein sport- und bewegungstherapeutisches Programm für Menschen mit psychischen Erkrankungen.

Zheng, B., Yu, C., Lin, L., Du, H., Lv, J., Guo, Y., Bian, Z., Chen, Y., Yu, M., Li, J., Chen, J., Chen, Z., Li, L., & The China Kadoorie Biobank Collaborative Group. (2017). Associations of domain-specific physical activities with insomnia symptoms among 0.5 million Chinese adults. *Journal of Sleep Research*, 26(3), 330–337.
<https://doi.org/10.1111/jsr.12507>